

April 2020 Vol. 113 No. 4

COMMENTARY

COVID-19: THE PANDEMIC OF OUR TIME	P46
[Read Article]	

EDITORIALS

IMAGING OF COVID-19; AN IRISH PERSPECTIVEP47 [Read Article]	7
PROVISION OF SERVICE DURING THE COVID-19 PANDEMIC	3
PREGNANCY AND SARS-COV-2: A NOVEL VIRUS IN A UNIQUE POPULATION)
PROTECTING OUR LONGEVITY DIVIDEND DURING COVID-19)
CARING FOR THE PSYCHOLOGICAL WELL-BEING OF FRONT LINE WORKERS DURING COVID-19P51 [Read Article]	Ĺ
SOCIAL MEDIA SPREAD DURING COVID-19: THE PROS AND CONS OF LIKES AND SHARESP52 [Read Article]	2

INTERVIEW

COVID-19 AND GENERAL PRACTICE: INTERVIEW WITH DR. RAY WALLEY, MEMBER OF THE NATIONAL	
COVID-19 GP LIAISON COMMITTEE.	P53
[Read Article]	

ORIGINAL PAPERS


> -Preview-45% 40% 35% 30% 25% Post-operative enoxaparin 20% Pre-operative enoxaparin Did not receive enoxaparin 15% 10% 5% 0% Post-operative **Pre-operative Did not receive** enoxaparin enoxaparin enoxaparin Figure 1: Distribution and timing of enoxaparin administration in breast surgery patients

Fullard et al report a 4% post-operative haematoma rate. The timing of LMWH did not affect the rate. [Read Article]

Costello et al found that among 110 cases of syncope, 62 had taken place at mass. The explanation is the multiple changes in posture. [Read Article]

-Preview				
Table 1: Times and durations of stance changes during mass.				
Position	Time			
Sit before Mass starts	Person dependent			
Stand	4.41 minutes			
Sit	3.26 minutes			
Stand	2.28minutes			
Sit	5.54 minutes			
Stand	3.54 minutes			
Sit	2.19 minutes			
Stand	51 seconds			
Kneel	3.59 minutes			
Stand	1.39 minutes			
Kneel	2.01 minutes			
Walk to Communion	Person dependent			
Kneel after communion	1.30 minutes			
Sit	2.18 minutes			
Stand	1.36 minutes			
Total	33.56 minutes approximately			

Grant and Harper found that in a series of 34 tonsillectomies and adeno-tonsillectomies, 6 children were admitted from the day ward or readmitted from home. [Read Article]



Laughlin et al describe the use of the neonatal sepsis calculator in the assessment of the risk of neonatal group B str. infection. The introduction of the calculator has reduced both cultures and antibiotic use. [Read Article]

Duraniar

-Preview- Table 1: Annual birth, numbers of blood cultures, antibiotic use within 24 hours and rate of blood culture draw.					
Year	2015	2016	2017	2018	January-March 2019
Total Babies (>500g)	9389	9037	8619	7923	2546
>37 weeks	8735	8428	7983	7310	2393
< 37 weeks	636	609	636	568	145
Missing GA	-	-	-	45	8
Number of blood cultures taken within first 72	1312	1149	1319	702	192
hours of life (All gestational ages)					
Antibiotics use within the first 24 hours of life	11.5%	10.4%	12.1%	6.7%	5.1%
Rate of Blood Culture Drawn / 1000 live births	140	127	153	89	76

OCCASIONAL PIECES

OPTIMISATION OF VITAMIN D STATUS FOR ENHANCED IMMUNO-PROTECTION AGAINST COVID-19......P58 [Read Article]

VITAMIN D, COVID-19 AND CHILDREN	P59
[Read Article]	

CASE REPORTS

A RARE CASE OF INFERTILITY: SRY POSITIVE 46, XX TESTICULAR DISORDER OF SEXUAL DIFFERENTIATION

Bogdanet et al describe a case of SRY positive 46 XX testicular disorder. The patient presented with a history of infertility.

[Read Article]

Table 1: Biochemical investigations.				
Blood Test	Value	Reference Range		
Testosterone	6.7 nmol/l	10.4-41.6 nmol/L		
LH	4.4 IU/L	1.8-12 IU/L		
FSH	43.1 IU/L	1-8 IU/L		
HbA1c	38 mmol/mol	20-40 mmol/mol		
IgA anti tTTG	<1 U/ml	0-10 U/ml		
PSA	0.74 ug/L	0-2.5 ug/L		
Free T4	13.2 pmol/L	10.5-22.0 pmol/L		
TSH	1.03 mIU/L	0.27-4.2 mIU/L		
Prolactin	112 mIU/L	86-324 mIU/L		

HEPATIC CYSTIC ECHINOCOCCOSIS (HYDATID CYST) IN A SIX YEAR OLDP61

Ali et al report a case of hydatid cyst in a 6 years old child. It was in the right lobe of the liver. It was treated with surgical removal and Albendazole and Praziquantal medication. [Read Article]



Figure 1: The T2 weighted MRI image of the liver in the coronal plane shows a single unilocular fluid containing cyst (36mm x 27mm x 51mm) in segment 6 of the liver, with an inner floating detached membrane, the so-called "water lily sign". No daughter cysts were identified.

-Preview-

LETTERS TO THE EDITOR

UMBILICAL CORD STRICTURE CAUSING INTRAUTERINE DEATH	P62
[Read Article]	

SHOULD ORAL SODIUM BICARBONATE BE USED TO PREVENT CHRONIC KIDNEY DISEASE PROGRESSION?...P65 [Read Article]

SUPPLY OF PERSONAL PROTECTIVE EQUIPMENT (PPE) DURING THE COVID-19 PANDEMIC......P66 [Read Article]

CONTINUAL PROFESSIONAL DEVELOPMENT

IMO Members can take our CPD Quiz to earn CPD/CME points.

CPD QUIZ

THE IRISH MEDICAL JOURNAL MAILING LIST

Subscribe below to receive the Irish Medical Journal newsletter. (There are 10 newsletters sent each year with new research updates.)

SUBSCRIBE