

Is high-concentration Nitrous oxide the safest option for short periods of procedural sedation in Paediatric Emergency Medicine

J. Coveney, U. Khan, T. Beattie

Department Child Life & Health, University of Edinburgh, Edinburgh, Scotland.

Dear Editor,

Since the 1990's there has been an increase in the number of invasive procedures being performed in paediatric emergency departments (PEDs) using procedural sedation and analgesia (PSA)^{1,2}. A recent survey of British and Irish PEDs found that 82% of departments performed PSA².

Timely PSA provides rapid pain relief and by allowing procedures to be completed in the PED reduces the need for hospital admission. While there are many benefits of PSA, as yet there is no perfect medication or combination of medications that provides PSA without any risk of adverse event.

Nitrous Oxide (N₂O) is a commonly used agent for PSA in PEDs. In general, N₂O at concentrations up to 50% provides minimal sedation, with a low reported incidence of severe adverse events (SAEs). At concentrations of 50% -70% it is more likely to generate moderate or deep sedation (with, in theory, greater potential for SAEs)

To inform PSA practice, we performed a literature review to identify the incidence of SAEs, significant interventions in response to SAEs and unplanned hospital admissions, occurring in children aged one to seventeen years, undergoing PSA using N₂O at concentrations between 51% and 70% in the PED setting.

Methodology from preferred reporting items for systematic reviews and meta-analyses statement (PRISMA) was followed. A comprehensive search of three electronic databases: Cochrane Central Register of Controlled Trials, Ovid MEDLINE and PubMed Central was completed.

SAEs were defined as: apnoea, laryngospasm, hypotension, bradycardia, complete airway obstruction, clinically apparent pulmonary aspiration, permanent neurologic injury, or death³. Significant interventions were defined as: positive pressure ventilation, endotracheal intubation, administration of vasoactive medication, administration of neuromuscular blockade or performance of chest compressions³.

An assessment of the risk of bias in included randomised controlled studies was performed using the of the Cochrane Bias Assessment Tool for randomised trials. An assessment of the risk of bias in included observational studies was performed using the Cochrane Risk of Bias in Non-randomized Studies tool.

Three RCTs (878 PSAs) and five observational studies (2,552 PSAs) were identified. The incidence of SAEs was 0.3 per 1,000 PSAs (1 episode of bradycardia which responded to the administration of oxygen). There was no reported case of intervention in response to SAEs. The incidence of unplanned hospital admission was 0.3/1,000 PSAs. The risk of bias assessment found two studies at low risk of bias, five studies at moderated risk of bias and one study at high risk of bias.

The incidence of SAEs identified is significantly lower than that reported with other commonly used PSA agents used in the PED (ketamine, propofol, midazolam, chloral hydrate)⁴.

Given the risk of bias assessment, a concern is that underreporting of SAEs in studies could lead to underestimation of the incidence of SAEs. However, the incidence of SAEs, interventions in response to SAEs and unplanned hospital admissions, were all low.

In summary, high-concentration N₂O is identified as having a lower incidence of SAEs than other PSA agents. The favourable safety profile should be considered when choosing a PSA agent in the PED.

Declarations of Conflicts of Interest:

None declared.

Corresponding author:

John Coveney,
Department Child Life & Health,
University of Edinburgh,
Edinburgh, Scotland.
E-Mail: johncoveney@gmail.com

References:

1. Borland M, Esson A, Babl F, Krieser D. Procedural sedation in children in the emergency department: a PREDICT study. *Emerg Med Australas* [Internet]. 2009 Feb

- [cited 2022 Sep 29];21(1):71–9. Available from:
<https://pubmed.ncbi.nlm.nih.gov/19254316/>
2. Vassallo J, Chernbumroong S, Malik N, Xu Y, Keene D, Gkoutos G, et al. 784 The landscape of paediatric procedural sedation in UK & Irish emergency departments; a PERUKI study. *Emerg Med J* [Internet]. 2022 Mar 1 [cited 2022 Nov 15];39(3):245–6. Available from: <https://emj.bmj.com/content/39/3/245.3>
 3. Bhatt M, Kennedy RM, Osmond MH, Krauss B, McAllister JD, Ansermino JM, et al. Consensus-based recommendations for standardizing terminology and reporting adverse events for emergency department procedural sedation and analgesia in children. *Ann Emerg Med* [Internet]. 2009 [cited 2022 Oct 8];53(4). Available from: <https://pubmed.ncbi.nlm.nih.gov/19026467/>
 4. Bhatt M, Johnson DW, Chan J, Taljaard M, Barrowman N, Farion KJ, et al. Risk Factors for Adverse Events in Emergency Department Procedural Sedation for Children. *JAMA Pediatr* [Internet]. 2017 Oct 1 [cited 2022 Nov 16];171(10):957–64. Available from: <https://pubmed.ncbi.nlm.nih.gov/28828486/>